

Providence College

DigitalCommons@Providence

Mathematics & Computer Science Student
Scholarship

Mathematics & Computer Science

4-22-2020

Simplexity of the n-cube

Peter Graziano

Follow this and additional works at: https://digitalcommons.providence.edu/computer_science_students



Part of the [Mathematics Commons](#)

Simplexity of the n -cube of Low Dimensions

Peter Graziano (Mathematics, '20)

Mathematics, Providence College

Objectives

- Recreate the preexisting results for the $n = 4$ case independently.
- Find a bound for the $n = 5$ case by a method besides brute force.
- Find a general pattern in the number of possible external faces of a simplex with respect to its possible volume.
- Find a general pattern in the minimal number of simplices for some n -cube.

Introduction

This project studied mathematical objects called n -cubes, which, like a square (2-cube) or a (3-)cube, have $\frac{n2^n}{2}$ edges such that all edges are the same length and any two edges which touch are perpendicular to each other in some plane, and their relation to mathematical objects called n -simplices, which, like triangles (2-simplices), have $n + 1$ vertices with no more than 3 in the same plane. Specifically, it studied how few simplices are needed to cover a n -cube for a particular value of n . The simplest case, when $n = 2$, simply shows how a square (2-cube) can be covered with two triangles (2-simplices), but any higher case has several different coverings possible. This project was concerned with finding a lower bound on the minimal number of simplices needed to cover the 4-cube and 5-cube.

The project was based upon a line of papers by Mara, Heiman, Cottle, Salee, and others using various techniques to tighten the bounds on optimal solutions for coverings (or triangulations) of the n -cube for particular values of n or for a general n .

Definitions


- An n -**simplex**, S^n , is an n -polytope with $n + 1$ vertices, $V = \{v_0, v_1, \dots, v_n\}$, such that $v_i - v_j$ is linearly independent, $\forall i, j \in V$.
- A **triangulation** of an n -polytope, P , is a finite set S of n -simplices such that S covers P completely and disjointly, or symbolically:
 - 1 $P = \cup S$
 - 2 For all $a, b \in S$, $a \cap b$ is a face of both a and b or $a \cap b = \emptyset$.

Process


In broad strokes, the process was as follows:

- Take simple geometric object, in this case, the n -cube
- Find a map to an algebraic object, in this case, the matrix representation of the n -cube
- Associate geometric quality with algebraic quantity, in this case, the volume of the simplices and the value of the determinant of a matrix which describes it.
- Use the known quantities to set up systems of inequalities.
- Solve the systems using linear programming algorithms

$n = 3$ case



(a) Trivial triangulation of the 3-cube



(b) Optimal triangulation of the 3-cube

Mathematical Section

In order to find the main result, a bound on the volume of a simplex was needed.

There are ways to represent a simplex as a square matrix C with only 1 or -1 in its entries. This allows the use of Hadamard's inequality (which concerns the maximum value of the determinant of a certain class of square matrices) to find a bound on the determinant of C , namely:

$$|\det C| \leq \frac{\binom{n}{2}}{2^{n-1}} \quad (1)$$

Since the volume of all the simplices in a triangulation of a cube is equal to the volume of the whole cube, a bound on the volume of a simplex leads to a bound on the number of possible simplices in a triangulation of a cube.

Main Result

Let T^n be a triangulation of I^n . Then,

$$(1) \min |T^3| \geq 5. \quad (2) \min |T^4| \geq 16. \quad (3) \min |T^5| \geq 60.$$

Conclusion

While doing the research for this project, I found that the actual optimal value of $|T^5| = 67$. The methods used to discover this, however, involved computing every possible 5-simplex and checking possible triangulations by computer. This project was the beginnings of an attempt to replicate this result by a more elegant method.

I found, however, that the method I used was not powerful enough to develop new independent results, and so more research would be needed.

References

- [1] Cottle, Richard W. "Minimal Triangulation of the 4-cube." *Discrete Mathematics* 40 pp.25-29, 1982.
- [2] Haiman, Mark. "A Simple and Relatively Efficient Triangulation of the n -Cube." *Discrete and Computational Geometry* 6 pp.287-289, 1991.
- [3] Mara, Patrick Scott. "Triangulations for the Cube." *Journal of Combinatorial Theory* 20 pp.170-177, 1976.
- [4] Sallee, John F. "A Triangulation of the n -cube." *Discrete Mathematics* 40 pp. 81-86, 1982.

Acknowledgements

I could not have done this project without the help of Dr. Su-Jeong Kang, to whom I am very grateful. I was funded through the Providence College Undergraduate Student Grant program.

Contact Information

- Email: pgrazian@friars.providence.edu