

5-27-2010

# Collisional Transfer of Population and Orientation in NaK

Seth T. Ashman

*Providence College, sashman@providence.edu*

C. M. Wolfe

*Lehigh University*

J. Huennekens

*Lehigh University*

B. Beser

*Temple University*

J. Bai

*Temple University*

*See next page for additional authors*

Follow this and additional works at: [http://digitalcommons.providence.edu/physics\\_fac](http://digitalcommons.providence.edu/physics_fac)



Part of the [Atomic, Molecular and Optical Physics Commons](#)

---

Ashman, Seth T.; Wolfe, C. M.; Huennekens, J.; Beser, B.; Bai, J.; and Lyyra, A. M., "Collisional Transfer of Population and Orientation in NaK" (2010). *Physics Faculty Publications*. 7.

[http://digitalcommons.providence.edu/physics\\_fac/7](http://digitalcommons.providence.edu/physics_fac/7)

This Conference Proceeding is brought to you for free and open access by the Physics at DigitalCommons@Providence. It has been accepted for inclusion in Physics Faculty Publications by an authorized administrator of DigitalCommons@Providence. For more information, please contact [mcaprio1@providence.edu](mailto:mcaprio1@providence.edu).

---

**Authors**

Seth T. Ashman, C. M. Wolfe, J. Huennekens, B. Beser, J. Bai, and A. M. Lyyra

# Collisional Transfer of Population and Orientation in NaK

C. M. Wolfe<sup>1</sup>, S. Ashman<sup>1</sup>, J. Huennekens<sup>1</sup>, B. Beser<sup>2</sup>, J. Bai<sup>2</sup>, A. M. Lyra<sup>2</sup>

<sup>1</sup>Lehigh University, <sup>2</sup>Temple University


Work Supported by the National Science Foundation



## Abstract

We report current work to study transfer of population and orientation in collisions of NaK molecules with argon and potassium atoms using polarization labeling (PL) and laser-induced fluorescence (LIF) spectroscopy. In the PL experiment, a circularly polarized pump laser excites a specific NaK  $A^1\Sigma^+(v'=16, J) \leftarrow X^1\Sigma^+(v''=0, J' \pm 1)$  transition, creating an orientation (non-uniform  $m_J$  level distribution) in both levels. The linearly polarized probe laser is scanned over various  $3^1\Pi(v, J' \pm 1) \leftarrow A^1\Sigma^+(v'=16, J)$  transitions. The probe laser passes through a crossed linear polarizer before detection, and signal is recorded if the probe laser polarization has been modified by the vapor (which occurs when it comes into resonance with an oriented level). Using both spectroscopic methods, analysis of weak collisional satellite lines adjacent to these directly populated lines, as a function of argon buffer gas pressure and cell temperature, allows us to discern separately the effects collisions with argon atoms and potassium atoms have on the population and orientation of the molecule. In addition, code has been written which provides a theoretical analysis of the process, through a solution of the density matrix equations of motion for the system.

## Background




- We use Optical-Optical Double Resonance (OODR) to study the NaK molecule in both Polarization Labeling and Laser-Induced Fluorescence spectroscopy
- Quantitative study of collisional population and orientation transfer
- Separately study the effects of collisions with Argon vs. Potassium


## Polarization Spectroscopy

Due to asymmetric  $m_J$  distribution created by circular polarized pump beam ( $\Delta m = +1$  for right circular polarization), vapor is birefringent for any probe that shares upper, lower, or both levels of pump transition

Note: linearly polarized probe can be thought of as equal components of left and right circular polarization. ( $\Delta m = -1$  and  $+1$ )




## Comparison Spectra




Comparison excitation scan showing both polarization spectroscopy (top) and laser-induced fluorescence spectroscopy (bottom)

Direct pump-probe transition is



Collisionally populated and oriented levels  $A^1\Sigma(16, 30 + \Delta J)$  are labeled by  $\Delta J$

## Collisional Transfer




Collisional effects on population and orientation can be divided into two types:

- Population and orientation are transferred from state 1 to state 2
- Population and orientation are lost from state 2 due to subsequent collisions and radiative decay

Empirical models have been developed to study population and orientation transfer, separating the effects of argon and potassium

## Transfer of Population (Fluorescence Spectroscopy)



Ratio of fluorescence intensities is roughly equal to the ratio of population densities in the collisionally and directly populated level

$$R_f = \frac{I'}{I} = \frac{I_2^F}{I_1^F} = \frac{n_2}{n_1}$$


Using a rate equation approximation, we model this ratio as

$$\frac{n_2}{n_1} = \frac{k_{Ar}^{\Delta J} n_{Ar} + k_K^{\Delta J} n_K}{\Gamma + k_{Ar}^Q n_{Ar} + k_K^Q n_K}$$

Here  $k^{\Delta J}$  is the rate of transfer of population from level 1 to the level characterized by  $\Delta J$  and  $k^Q$  is the total collisional quenching rate

Intensity ratios are fitted as a function of argon and potassium density in order to obtain population transfer rates

## Transfer of Orientation (Polarization Spectroscopy)



Polarization signal depends on the difference of absorption between left and right circular polarization components of the linearly polarized probe beam

Can model this using: Signal  $\propto \sum \text{Right Absorptions} - \sum \text{Left Absorptions}$

$$= \sum_{m=-J}^{+J} n_m F_m^R - \sum_{m=-J}^{+J} n_m F_m^L = \sum_{m=-J}^{+J} n_m (F_m^R - F_m^L) = f(J, J') \sum_{m=-J}^{+J} m n_m = f(J, J') \langle m \rangle \sum_{m=-J}^{+J} n_m$$

In this expression,  $F$  factors are relative absorption coefficients:

$$| \langle \beta_1, J_1, m_1 | \vec{r} | \beta_2, J_2, m_2 \rangle |^2 = F_{J_1 J_2}^{m_1 m_2} | \langle \beta_1 || r || \beta_2 \rangle |^2$$

Evaluation of difference in  $F$  factors yields  $F_m^R - F_m^L = f(J, J') m$

Thus the ratio of polarization spectroscopy signal strengths for the collisional and direct line are roughly equal to the ratio of orientation multiplied by the population of the levels. We can use a rate equation model to solve for the ratio of the intensities of the collisional and direct line:

$$R_p = \frac{\langle m_2 \rangle n_2}{\langle m_1 \rangle n_1} = \frac{k_{Ar}^{\Delta J, O} n_{Ar} + k_K^{\Delta J, O} n_K}{1 + \frac{g_{Ar}}{\Gamma} n_{Ar} + \frac{g_K}{\Gamma} n_K}$$

Here  $k^{\Delta J, O}$  is the rate of transfer of orientation from level 1 to the level characterized by  $\Delta J$ , and  $g$  is the total collisional rate of destruction of orientation. Separating out the population and orientation dependence yields

$$k_{Ar}^{\Delta J, O} = k_{Ar}^{\Delta J} (1 - f_{Ar}^{\Delta J}) \quad g_{Ar} = k_{Ar}^Q + g'_{Ar}$$

$$k_K^{\Delta J, O} = k_K^{\Delta J} (1 - f_K^{\Delta J}) \quad g_K = k_K^Q + g'_K$$

$f_{Ar, K}$  represents the fraction of orientation destroyed in the initial collision with either Ar or K, and  $g'$  is the rate of destruction of orientation by subsequent collisions that do not change  $J$ .  $g'$  is small compared to  $k^Q$ , so we approximate


$$g'_{Ar} = \frac{1}{2} (f_{Ar}^{-1} k_{Ar}^{-1} + f_{Ar}^{+1} k_{Ar}^{+1})$$

$$g'_K = \frac{1}{2} (f_K^{-1} k_K^{-1} + f_K^{+1} k_K^{+1})$$

A fit of intensity ratios obtained from fluorescence and polarization spectroscopy, using our empirical formulas  $R_f$  and  $R_p$ , as functions of Ar or K number density has been carried out, for  $\Delta J = \pm 1, \pm 2, \pm 3, \pm 4$

## Fitting Results

The parameters  $k_{Ar}$ ,  $k_K$ ,  $f_{Ar}$ , and  $f_K$  give us information on the population transferred, as well as the orientation destroyed, in the initial collision from the directly populated intermediate level to the collisional level.



- Only  $\Delta J = \pm 2, \pm 4, \dots$  transitions observed in homonuclear molecules (like Rb<sub>2</sub>)
- $k_{Ar}$  values are much larger for even numbered collisional transitions than for odd numbered transitions
- $k_K$  values do not seem to show such a propensity
- $f$  values show that collisions with potassium atoms are more likely to destroy angular momentum orientation than collisions with argon atoms

$$k_{Ar}^Q = 1.16 \pm .07 \times 10^{-9} \text{ cm}^3 \text{ s}^{-1}$$


$$k_K^Q = 5.3 \pm .6 \times 10^{-9} \text{ cm}^3 \text{ s}^{-1}$$

## Density Matrix Model

To completely describe a system of quantum mechanical energy levels in a statistical manner, it is necessary to solve the density matrix equations of motion

$$\dot{\rho}_{nm} = -\frac{i}{\hbar} \sum_k (H_{nk} \rho_{km} - \rho_{nk} H_{km}) + \text{relaxation terms}$$

A diagonal element  $\rho_{nn}$  represents the population of sublevel  $n$ , while an off-diagonal element  $\rho_{nm}$  represents the laser-induced coherence between sublevels  $n$  and  $m$ . A computer code has been written which solves the density matrix equations of motion for either three coupled angular momentum states for the direct probe model, or four states for the collisional model.




Model Includes

- Laser coherence terms
- Fluorescence into and out of various levels
- Transit relaxation
- Additional "dump" levels representing other ground/intermediate levels
- Velocity group and laser detuning loops

Collisional model is equipped to incorporate any theoretical model for collisional transfer of population and orientation linking states "2" and "c"

## Sample Computer Model Results


### Detuning Loop Calculations



High Probe Power

Low Probe Power

### Intermediate state orientations predicted for various pump-probe power regimes



Low Pump, Zero Probe

Medium Pump, Zero Probe

High Pump, Zero Probe

Medium Pump, Low Probe

Medium Pump, Medium Probe

Medium Pump, High Probe

## Future Plans

- Find an appropriate theoretical model for the change in orientation during a collision to incorporate into density matrix
- Investigate ground state collisional lines
  - Thus far we have only investigated excited state collisional lines
  - $v$ -changing collisions have been observed in ground state. We plan to investigate these in greater detail
- At Temple U., we have observed ground state collisional lines for very large  $\Delta J$  (up to  $\Delta J = 58$ ) in polarization spectroscopy of Rb<sub>2</sub>
  - How can orientation be preserved in collisions where  $J$  changes by a very large amount?
- Effects on orientation/population by other collision partners (He, Xe, Kr)
- Propensity toward  $\Delta J = \pm 2, \pm 4$  not observed in NaCs
  - Is this due to NaCs being "more heteronuclear" than NaK?