

Providence College

DigitalCommons@Providence

---

Biology Student Scholarship

Biology

---

Spring 4-29-2021

## Elbow Motion in Walking Alligators

Thao Pham

*Providence College*

Luciana Emmanuelli

*Providence College*

Megan Rowlings

*Providence College*

Nina Pitre

*Providence College*

Follow this and additional works at: [https://digitalcommons.providence.edu/bio\\_students](https://digitalcommons.providence.edu/bio_students)



Part of the [Biology Commons](#)

---


Pham, Thao; Emmanuelli, Luciana; Rowlings, Megan; and Pitre, Nina, "Elbow Motion in Walking Alligators" (2021). *Biology Student Scholarship*. 17.

[https://digitalcommons.providence.edu/bio\\_students/17](https://digitalcommons.providence.edu/bio_students/17)

This Poster is brought to you for free and open access by the Biology at DigitalCommons@Providence. It has been accepted for inclusion in Biology Student Scholarship by an authorized administrator of DigitalCommons@Providence. For more information, please contact [dps@providence.edu](mailto:dps@providence.edu).

## Introduction:

Crocodylia regularly use a "high-walk" during terrestrial locomotion unlike other non-mammals, thus drawing interest from both biomechanical and evolutionary research communities. Here, we employ the highly accurate XROMM (X-ray Reconstruction of Moving Morphology) method to measure joint motion between the three forelimb bones (humerus, ulna, and radius) of alligators performing high-walks on a treadmill. This method literally "re-animates" bone motion by combining bi-planar X-ray video with digital bone models from CT scans in the animation program Maya (Autodesk).



## Anatomical Model:




Figure 1. Alligator elbow joint. The elbow consists of articulations between three bones forming the **humero-ulnar joint**, **humero-radial joint**, and **radio-ulnar joint**. In this study, we limit our focus to the **humero-ulnar joint**. H = humerus, U = ulna, R = radius.

## Hypothesis:

We predict that there will be a larger range of motion for flexion/extension in both the humero-ulnar and humero-radial joint, while the radio-ulnar joint will show a significantly smaller range. We also predict that the radio-ulnar joint will show a larger range of motion for long-axis rotation than the other two joints. For range of translation, we predict that none of the three joints will show a significant range of motion.

## Methods:

We used XROMM (Brainerd et al. 2010, Knorlein et al. 2016) "re-animate" 3D skeletal motion by combining in vivo x-ray videos with Computed Tomography (CT Scans). We calculated relative joint motion for the humero-radial, humero-ulnar, and radio-ulnar joints, then scaled the animation into half-strides to compare ranges of joint motion for 3 rotational and 3 translational degrees of freedom.

# Elbow Motion in Walking Alligators

Thao Pham, Megan Rowlings, Nina Pitre, Luciana Emmanuelli, and David Baier - Providence College

## Elbow Motion:




Figure 2. Humero-ulnar joint motion across three stances and three swing phases. Animation was scaled from 0-100 frames for each half stride.

Orange = long-axis rotation (x-axis), Green = Abduction/Adduction (y-axis), and Blue = Flexion/Extension (z-axis)




Figure 3. Humero-radial joint motion across three stances and three swing phases. Animation was scaled from 0-100 frames for each half stride.

Orange = long-axis rotation (x-axis), Green = Abduction/Adduction (y-axis) and Blue = Flexion/Extension (z-axis)




Figure 4. Radio-ulnar joint motion across three stances and three swing phases. Animation was scaled from 0-100 frames for each half stride.

Orange = long-axis rotation, (x-axis) Green = Abduction/Adduction (y-axis), and Blue = Flexion/Extension (z-axis)

## Conclusions:

From the joint motion data graphed, we observed that the humero-ulnar and humero-radial joint showed a larger range of motion for flexion/extension, and a smaller range of motion for both long axis rotation and abduction/adduction. Range of translation is also small for both joints. We also expected the radio-ulnar joint to show a large range of motion for long-axis rotation and a smaller range of motion for flexion/extension and abduction/adduction, however we did not expect that there would be a large range of translation from the radio-ulnar joint.

## Future Research:

In the future, it may be beneficial to look at multiple alligators to have a broader scope of the rotational degree measurements.

## References:

- Baier et al. (2013) Three-dimensional skeletal kinematics of the shoulder girdle and forelimb in walking Alligator. *J. Anat.*(223)
- Baier et al. (2013) Three-Dimensional, High-Resolution Skeletal Kinematics of the Avian Wing and Shoulder during Ascending Flapping Flight and Uphill Flap-Running. *PLoS ONE*.
- Brainerd et al. (2010) X-ray reconstruction of moving morphology (XROMM): precision, accuracy and applications in comparative biomechanics research. *J. Zool*
- Knorlein et al. (2016) Validation of XMALab software for marker-based XROMM. *J. Exp. Biol.*